

Membrane and Noncommutativity

Rabin Banerjee*,

Institute of Particle and Nuclear Studies,
High Energy Accelerator Research Organisation (KEK),
Tsukuba 305-0801, Japan

Biswajit Chakraborty[†] and Kuldeep Kumar[‡]

S. N. Bose National Centre for Basic Sciences,
Block JD, Sector III, Salt Lake, Kolkata-700 098, India

February 1, 2008

Abstract

We analyse the dynamics of an open membrane, both for the free case and when it is coupled to a background three-form, whose boundary is attached to p -branes. The role of boundary conditions and constraints in the Nambu-Goto and Polyakov formulations is studied. The low-energy approximation that effectively reduces the membrane to an open string is examined in detail. Noncommutative features of the boundary string coordinates, where the cylindrical membrane is attached to the Dp -branes, are revealed by algebraic consistency arguments and not by treating boundary conditions as primary constraints, as is usually done. The exact form of the noncommutative algebra is obtained in the low-energy limit.

Keywords: Constrained hamiltonian analysis, Membranes, Noncommutativity

PACS: 11.10.Nx, 11.25.-w, 11.25.-q, 11.10.Ef

1 Introduction

Over the last decade string theory has been gradually replaced by M-theory as the most natural candidate for a fundamental description of nature. While a complete definition of M-theory is yet to be given, it is believed that the five perturbatively consistent string theories are different phases of this theory. With the replacement of string theory by M-theory, the string itself has lost its position as the main candidate for the fundamental degree of freedom. Instead, higher-dimensional extended objects like membranes are being considered [1]. Indeed it is known that membrane and five-brane occur naturally in eleven-dimensional supergravity, which is argued to be the low-energy limit of M-theory. Also, string theory is effectively described by the low-energy dynamics of a system of

*On leave of absence from S. N. Bose National Centre for Basic Sciences, Kolkata, India;
rabin@post.kek.jp, rabin@bose.res.in

[†]biswajit@bose.res.in

[‡]kuldeep@bose.res.in

branes. For instance, the membrane of M-theory may be “wrapped” around the compact direction of radius R to become the fundamental string of type-IIA string theory, in the limit of vanishing radius.

An intriguing connection between string theory, noncommutative geometry, noncommutative (as well as ordinary) Yang-Mills theory was revealed in [2]. With the shift in focus from string theory to M-theory, there has been a flurry of activity in analysing noncommutativity in membranes, specifically when an open membrane that couples to a three-form, ends on a D-brane [3, 6, 7, 8]. The motivation of the present paper is to further this investigation, but with a new perspective and methodology, as explained below.

The study of noncommutative properties in membranes is more involved than the analogous study in the string case since the equations to be solved are nonlinear. Naturally, in contrast to the string situation, the results could be obtained only under some approximations. It is useful to recapitulate how noncommutativity is derived in either the string coupled to the two-form or the membrane coupled to the three-form. There are non-trivial boundary conditions which are incompatible with the basic Poisson brackets of the theory. These boundary conditions are considered as primary constraints in the algorithm of Dirac’s constrained hamiltonian dynamics [4, 5, 6, 7, 8, 9, 10]. The primary constraints lead to secondary constraints. Noncommutativity is manifested through the occurrence of non-trivial Dirac brackets. The brackets are found to be gauge dependent, but there is no gauge where it can be made to vanish.

Recently, an alternative approach to deal with noncommutativity in strings was advocated in a paper [11] involving two of us. Contrary to other approaches, the boundary conditions are not interpreted as primary constraints. The noncommutative algebra emerges from a set of consistency requirements. It is rather similar in spirit to the original analysis of [12] where a modified algebra, involving the periodic delta function instead of the usual one, was found for the coordinates and their conjugate momenta, in the example of the free Nambu-Goto (NG) string.

In this paper we adopt our previous strategy for strings to the membrane model. We discuss both the NG and Polyakov forms of action, although noncommutativity is explicitly considered only in the latter formulation. The similarities or otherwise in the analysis of the two actions are illuminated. Analogous to the set of orthonormal gauge fixing conditions given for the free Nambu-Goto string [11, 12], we derive a set of quasi-orthonormal gauge conditions for the free NG membrane. Just as the orthonormal gauge in the NG string corresponds to the conformal gauge in the Polyakov string, we find out the analogue of the quasi-orthonormal gauge in the Polyakov membrane. It corresponds to a choice of the metric that leads to equations of motion that can be explicitly solved in the light-front coordinates [13]. The structure and implications of the boundary conditions in the two formulations have been elaborated. In the NG case, the conditions involve the velocities that cannot be inverted so that a phase-space formulation is problematic. Only by fixing a gauge is it possible to get hold of a phase-space description. In the Polyakov type, on the other hand, the boundary condition is expressible in phase-space variables without the need of any gauge choice. This is because the metric itself is regarded as an independent field. In this sense, therefore, there is no qualitative difference between string and membrane boundary conditions, since even in the NG string, a gauge fixing is required for writing the boundary conditions in terms of phase-space variables. We thus differ from [7] where it is claimed that it is imperative in the membrane case, as opposed

to the string case, to gauge fix in order to express the boundary conditions in phase-space coordinates as a first step in the hamiltonian formalism.

The mandatory gauge fixing in the NG membrane, as we shall show, converts the reparametrization-invariant (first-class) system into a second-class one, necessitating the use of Dirac brackets. This involves the inversion of highly non-linear expressions, so that approximations become essential to make any progress. Hence we avoid this formulation in favour of the Polyakov version, where gauge fixing is not mandatory.

A detailed constrained hamiltonian analysis of the free bosonic Polyakov membrane naturally leads to three restrictions on the world-volume metric. These are found to be identical to those obtained by counting the independent degrees of freedom. Unlike the case of the classical string where there are three components of the metric and three continuous symmetries (two diffeomorphism symmetries and one scale symmetry), leading to a complete specification of the metric by gauge fixing, for the membrane there are six independent metric components and only three diffeomorphism symmetries. Thus only three restrictions on the metric can be imposed. Interestingly, the restrictions usually put in by hand [13] to perform calculations in the light-front coordinates are obtained directly in our hamiltonian formalism. This gauge fixing is only partial in the sense that the non-trivial gauge generating first-class constraints remain unaffected. Effectively, therefore, it is a gauge-independent hamiltonian formalism. We show that the boundary string coordinates corresponding to the membrane-D p -brane system (i.e. when the boundary of the open membrane is attached to p -branes) satisfy the usual Poisson algebra without any noncommutativity. By imposing further gauge conditions, it is possible to simulate a situation where the cylindrical membrane is wrapped around a circle of vanishing radius so that the open membrane passes over to an open string. The boundary conditions of the membrane reduce to the well-known Neumann boundary conditions of the string in the conformal gauge, just as the membrane metric reduces to the conformal metric of the Polyakov string.

Next, the interacting membrane in the presence of a constant three-form tensor potential is discussed. Proceeding in a gauge-independent manner, it is shown that, contrary to the free theory, the boundary string coordinates must be noncommutative. This is shown from certain algebraic conditions. However, in contrast to the string case where it was possible to solve these equations [11], here an explicit solution is prevented from the non-linear structure. Nevertheless, by passing to the low-energy limit (wrapping the membrane on a circle of vanishingly small radius), the explicit form of the noncommutativity in an open string, whose end points are attached to a D-brane, are reproduced.

The paper is organised as follows: In section **2**, the free NG membrane is discussed and the form of the quasi-orthonormal gauge conditions, which act as the analogue of the orthonormal gauge conditions in the NG string [12], is derived. The role of the boundary conditions in maintaining stability of the membrane is discussed. The free Polyakov membrane is considered in section **3**, where its detailed constrained hamiltonian account is given. The complete form of the energy-momentum tensor is derived. All components of this tensor are written as a linear combination of the constraints. This is a generalization of the string case since even though Weyl symmetry is absent in the membrane, the energy-momentum tensor has a (weakly) vanishing trace; namely, it vanishes only on the constraint shell. The brackets for the free theory with a cylindrical topology for the membrane, computed in section **4**, yield the expected Poisson algebra without any noncommutativity. The low-energy limit where the membrane is approximated by the

string, is discussed in section 5. Section 6 gives an analysis of the interacting theory. General algebraic requirements enforce a noncommutativity of the boundary coordinates of the membrane, which are attached to the p -branes. No gauge fixing or approximation is needed to reveal this noncommutativity. The explicit structure of the algebra is once again computed in the low-energy approximation, when the result agrees with the conformal-gauge expression for the noncommutativity among the coordinates of the end points of the string attached to D-branes. Concluding remarks are given in section 7. An appendix, summarizing the basic results of our earlier paper [11] on strings, has been included for easy comparison with the membrane analysis.

2 The Free Nambu-Goto Membrane

A dynamical membrane moving in $D-1$ spatial dimensions sweeps out a three-dimensional world-volume in D -dimensional space-time. We use a metric with signature $(-, +, +, \dots, +)$ in the target space whose indices are $\mu, \nu = 0, 1, 2, \dots, (D-1)$. We can locally choose a set of three coordinates $\sigma^i, i = 0, 1, 2$, on the world-volume to parameterize it. We shall sometime use the notation $\tau = \sigma^0$ and the indices $a, b, \dots$ to describe “spatial” coordinates $\sigma^a, a = 1, 2$, on the membrane world-volume. In such a coordinate system, the motion of the membrane through space-time is described by a set of D functions $X^\mu(\sigma^0, \sigma^1, \sigma^2)$ which are the membrane coordinates in the target space.

Although we are going to study the noncommutativity through the Polyakov action, we find it convenient to briefly discuss the NG action also. The NG analysis will be just an extension of the string case, considered in [12]. The NG action for a membrane moving in flat space-time is given by the integrated proper volume swept out by the membrane:

$$S_{NG} = -T \int_{\Sigma} d^3\sigma \sqrt{-h} \equiv \int_{\Sigma} d^3\sigma \mathcal{L}_{NG}(X^\mu, \partial_i X^\mu), \quad (1)$$

where T is a constant which can be interpreted as the membrane tension and $h = \det h_{ij}$ with

$$h_{ij} = \partial_i X^\mu \partial_j X_\mu \quad (2)$$

being the induced metric on (2+1)-dimensional world-volume, which is nothing but the pullback of the flat space-time metric on this three-dimensional sub-manifold. This induced metric, however, does not have the status of an independent field in the world-volume; it is rather determined through the embedding fields X^μ . The Lagrangian density is $\mathcal{L}_{NG} = -T\sqrt{-h}$. The Euler-Lagrange equation is given by

$$\partial_i (\sqrt{-h} h^{ij} \partial_j X^\mu) = 0, \quad (3)$$

while the boundary conditions are given by

$$\mathcal{P}_\mu^a \Big|_{\partial\Sigma} = -T\sqrt{-h} \partial^a X_\mu \Big|_{\partial\Sigma} = 0, \quad (4)$$

where

$$\mathcal{P}_\mu^i = \frac{\partial \mathcal{L}_{NG}}{\partial(\partial_i X^\mu)} = -T\sqrt{-h} \partial^i X_\mu \quad (5)$$

and $\partial\Sigma$ represents the boundary. The components $\mathcal{P}_\mu^0 \equiv \Pi_\mu$ are the canonical momenta conjugate to X^μ . Using this, the Euler-Lagrange equation (3) can be rewritten as

$$\partial_0 \Pi^\mu + \partial_a \mathcal{P}^{a\mu} = 0. \quad (6)$$

It can be seen easily that the theory admits the following primary constraints:

$$\psi \equiv \Pi^2 + T^2 \bar{h} \approx 0, \quad (7)$$

$$\phi_a \equiv \Pi_\mu \partial_a X^\mu \approx 0, \quad a = 1, 2, \quad (8)$$

where $\Pi^2 \equiv \Pi^\mu \Pi_\mu$ and $\bar{h} = \det h_{ab} = h_{11}h_{22} - (h_{12})^2$. These constraints are first-class since the brackets between them vanish weakly:

$$\begin{aligned} \{\psi(\tau, \vec{\sigma}), \psi(\tau, \vec{\sigma}')\} &= 4T^2 [\{h_{22}(\tau, \vec{\sigma}) \partial_1 \delta(\vec{\sigma} - \vec{\sigma}') - h_{12}(\tau, \vec{\sigma}) \partial_2 \delta(\vec{\sigma} - \vec{\sigma}')\} \phi_1(\tau, \vec{\sigma}) \\ &\quad + \{h_{11}(\tau, \vec{\sigma}) \partial_2 \delta(\vec{\sigma} - \vec{\sigma}') - h_{12}(\tau, \vec{\sigma}) \partial_1 \delta(\vec{\sigma} - \vec{\sigma}')\} \phi_2(\tau, \vec{\sigma}) \\ &\quad - \{h_{22}(\tau, \vec{\sigma}') \partial'_1 \delta(\vec{\sigma} - \vec{\sigma}') - h_{12}(\tau, \vec{\sigma}') \partial'_2 \delta(\vec{\sigma} - \vec{\sigma}')\} \phi_1(\tau, \vec{\sigma}') \\ &\quad - \{h_{11}(\tau, \vec{\sigma}') \partial'_2 \delta(\vec{\sigma} - \vec{\sigma}') - h_{12}(\tau, \vec{\sigma}') \partial'_1 \delta(\vec{\sigma} - \vec{\sigma}')\} \phi_2(\tau, \vec{\sigma}')] \\ &\approx 0, \end{aligned} \quad (9)$$

$$\{\phi_a(\tau, \vec{\sigma}), \phi_b(\tau, \vec{\sigma}')\} = \phi_b(\tau, \vec{\sigma}) \partial_a \delta(\vec{\sigma} - \vec{\sigma}') - \phi_a(\tau, \vec{\sigma}') \partial'_b \delta(\vec{\sigma} - \vec{\sigma}') \approx 0,$$

$$\{\psi(\tau, \vec{\sigma}), \phi_a(\tau, \vec{\sigma}')\} = 2\psi(\tau, \vec{\sigma}) \partial_a \delta(\vec{\sigma} - \vec{\sigma}') + \partial_a \psi \delta(\vec{\sigma} - \vec{\sigma}') \approx 0,$$

where $\partial'_a = \frac{\partial}{\partial \sigma'^a}$ and $\vec{\sigma} = (\sigma^1, \sigma^2)$.

The canonical world-volume energy-momentum tensor *density*¹ can be obtained through Noether theorem:

$$[\theta_C]^i_j = \frac{\partial \mathcal{L}_{NG}}{\partial (\partial_i X^\mu)} \partial_j X^\mu - \delta^i_j \mathcal{L}_{NG}. \quad (10)$$

In particular, $[\theta_C]^0_0 = 0$, $[\theta_C]^0_a = \phi_a \approx 0$, $[\theta_C]^a_0 = 0$ and $[\theta_C]^a_b = 0$. We notice that the canonical Hamiltonian density, $\mathcal{H}_C = [\theta_C]^0_0$, obtained by Legendre transformation, vanishes strongly. Since the canonical energy-momentum tensor density is first-class, we may add to it a linear combination of first-class constraints with tensor-valued coefficients to write down the total energy-momentum tensor density as

$$\theta^i_j = U^i_j \psi + V^{ai}_j \phi_a \approx 0. \quad (11)$$

The generators of τ and σ^a translations are

$$H_T = \int d^2 \sigma \theta^0_0, \quad H_a = \int d^2 \sigma \theta^0_a. \quad (12)$$

As one can easily see, there are no secondary constraints. The Hamilton's equation $\dot{X}^\mu = \{X^\mu, H_T\}$ gives

$$\partial_0 X^\mu = 2U^0_0 \Pi^\mu + V^{a0}_0 \partial_a X^\mu,$$

which reproduces the definition of momenta Π^μ for the following choice of U^0_0 and V^{a0}_0 :

$$U^0_0 = \frac{\sqrt{-h}}{2T\bar{h}}, \quad V^{a0}_0 = -\frac{h h^{0a}}{\bar{h}} = \bar{h}^{ab} h_{0b}, \quad (13)$$

where $\bar{h}^{ab}$ ($\neq h^{ab}$, which is obtained by chopping off first row and first column from h^{ij} , the inverse of h_{ij}) is the inverse of h_{ab} in the two-dimensional subspace. The other equation, $\dot{\Pi}^\mu = \{\Pi^\mu, H_T\}$, reproduces the Euler-Lagrange equation (3) whereas $\partial_a X^\mu = \{X^\mu, H_a\}$ gives

$$\partial_a X^\mu = 2U^0_a \Pi^\mu + V^{b0}_a \partial_b X^\mu,$$

¹Note that $\mathcal{L}_{NG}$ transforms as a scalar *density* under diffeomorphism.

which is satisfied for

$$U^0_a = 0, \quad V^{b0}_a = \delta^b_a. \quad (14)$$

Coming to the conserved Poincaré generators in the target space, the translational generator is given by

$$P_\mu = \int d^2\sigma \Pi_\mu,$$

and the angular momentum generator is given by

$$M^{\mu\nu} = \int d^2\sigma (X^\mu \Pi^\nu - X^\nu \Pi^\mu).$$

As can be easily checked, these generators generate appropriate Poincaré transformations. The above analysis can be generalized in a straightforward manner to an arbitrary p -brane.

There is an interesting implication of the boundary conditions (4). For a cylindrical membrane with $\sigma^1 \in [0, \pi]$, $\sigma^2 \in [0, 2\pi]$, σ^2 representing the compact direction, the boundary condition is written as

$$\mathcal{P}^1_\mu|_{\sigma^1=0,\pi} = -T\sqrt{-h}\partial^1 X_\mu|_{\sigma^1=0,\pi} = 0.$$

Squaring the above equation, we get

$$hh^{11}|_{\sigma^1=0,\pi} = [h_{00}h_{22} - (h_{02})^2]|_{\sigma^1=0,\pi} = 0, \quad (15)$$

which implies

$$h_{00}|_{\sigma^1=0,\pi} = \frac{(h_{02})^2}{h_{22}}|_{\sigma^1=0,\pi}. \quad (16)$$

However, h_{22} is strictly positive and cannot vanish at the boundary in order to prevent it from collapsing to a point as the length of the boundary is given by $\int_0^{2\pi} \sqrt{h_{22}} d\sigma^2$. This indicates that

$$\dot{X}^2|_{\sigma^1=0,\pi} = h_{00}|_{\sigma^1=0,\pi} \geq 0$$

so that the points on the boundary move along either a space-like or light-like trajectory. If we now demand that the speed of these boundary points should not exceed the speed of light then we must have $h_{02}|_{\sigma^1=0,\pi} = 0$ in Eq. (16) so that

$$\dot{X}^2|_{\sigma^1=0,\pi} = h_{00}|_{\sigma^1=0,\pi} = 0.$$

Therefore the boundary points move with the speed of light which is a direct generalization of the string case where a similar result holds. For a square membrane with $\sigma^1, \sigma^2 \in [0, \pi]$, the boundary conditions (4) are written as

$$\begin{aligned} \mathcal{P}^1_\mu|_{\sigma^1=0,\pi} &= -T\sqrt{-h}\partial^1 X_\mu|_{\sigma^1=0,\pi} = 0, \\ \mathcal{P}^2_\mu|_{\sigma^2=0,\pi} &= -T\sqrt{-h}\partial^2 X_\mu|_{\sigma^2=0,\pi} = 0. \end{aligned}$$

Therefore, in addition to Eq. (15), we also have

$$hh^{22}|_{\sigma^2=0,\pi} = [h_{00}h_{11} - (h_{01})^2]|_{\sigma^2=0,\pi} = 0.$$

Proceeding just as in the case of cylindrical membrane, we find that we must have $h_{02}|_{\sigma^1=0,\pi} = 0$ and $h_{01}|_{\sigma^2=0,\pi} = 0$ so that

$$\dot{X}^2|_{\sigma^1=0,\pi} = 0 = \dot{X}^2|_{\sigma^2=0,\pi},$$

which shows that the boundary points move with the speed of light. Also, since $h_{0a} \approx 0$ at the boundary, for both the cylindrical or square topology, it implies that the vector $\partial_0 X^\mu$ is not only null, but also orthogonal to all directions tangent to the membrane world-volume. Hence the boundary points move with the speed of light, perpendicularly to the membrane. This peculiar motion is exactly reminiscent of the string case. The tension in the free membrane would cause it to collapse. This is prevented by the angular momentum generated by the boundary motion, just as the collapse of the free string is thwarted by a similar motion of the string end points [14].

Quasi-orthonormal gauge fixing conditions

As we shall see now, the membrane case, or any p -brane with $p > 1$ for that matter, involves some subtle issues. The first step is to provide a set of complete gauge fixing conditions. Taking a cue from the previous analysis we would like to generalize the condition $h_{0a} \approx 0$, so that it holds everywhere, instead of just at the boundary. This is also quite similar in spirit to what is done for implementing the orthonormal gauge in the string case. Indeed, following the string analysis of [12], we first impose the following gauge fixing conditions:

$$\lambda_\mu \left(X^\mu(\tau, \vec{\sigma}) - \frac{P^\mu \tau}{TA} \right) \approx 0, \quad (17)$$

$$\lambda_\mu \left(\Pi^\mu(\tau, \vec{\sigma}) - \frac{P^\mu}{A} \right) \approx 0, \quad (18)$$

where λ_μ is an arbitrary constant vector and A is taken to be the “parametric area” of the membrane. For example, if the membrane is of square topology with $\sigma^1, \sigma^2 \in [0, \pi]$, it will be π^2 and for cylindrical topology with $\sigma^1 \in [0, \pi], \sigma^2 \in [0, 2\pi]$ (membrane periodic along σ^2 -direction), it will be $2\pi^2$. Clearly, this “parametric area” is not an invariant quantity under two-dimensional diffeomorphism. One can think of the square or cylindrical membrane to be flat at one instant to admit a Cartesian-like coordinate system on the membrane surface which will provide a coordinate chart for it during its future time evolution.

Differentiating Eq. (17) with respect to τ and using Eq. (18), we get

$$\lambda \cdot \dot{X} \approx \frac{\lambda \cdot P}{TA} \approx \frac{\lambda \cdot \Pi}{T}. \quad (19)$$

Differentiating Eq. (17) with respect to σ^a , and Eq. (18) with respect to τ we get

$$\lambda \cdot \partial_a X \approx 0, \quad (20)$$

$$\partial_0 (\lambda \cdot \Pi) \approx 0. \quad (21)$$

Using Eq. (21), it follows from the form (6) of Euler-Lagrange equation that

$$\partial_a (\lambda \cdot \mathcal{P}^a) \approx 0. \quad (22)$$

Upon contraction with λ^μ , the boundary conditions (4) give

$$\lambda \cdot \mathcal{P}^a|_{\partial\Sigma} = 0. \quad (23)$$

Now we impose an additional gauge fixing condition²

$$\varepsilon^{ab}\partial_a(\lambda \cdot \mathcal{P}_b) \approx 0. \quad (24)$$

Thus, we have from Eqs. (22) and (24) both the divergence and curl vanishing for the vector field $(\lambda \cdot \mathcal{P}^a)$ in the 2-dimensional membrane, which is also subjected to the boundary conditions (23). We thus have

$$\lambda \cdot \mathcal{P}^a = 0 \quad \forall \sigma^a. \quad (25)$$

In view of Eq. (20), we have

$$\lambda \cdot \Pi \approx -T\sqrt{-h}h^{00}(\lambda \cdot \partial_0 X),$$

which, using Eq. (19) gives

$$h^{00}\sqrt{-h} \approx -1. \quad (26)$$

Using Eqs. (19) and (20), Eq. (25) gives $h^{0a} \approx 0$ which implies

$$h_{0a} \approx 0, \quad h^{00} \approx \frac{1}{h_{00}}. \quad (27)$$

From Eqs. (26) and (27) it follows that

$$h_{00} + \bar{h} \approx 0. \quad (28)$$

Observe that the term quasi-orthonormality in this case means that the time-like vector ∂_0 is orthogonal to the space-like vectors ∂_a , which follows from Eq. (27). However, the two space-like directions ∂_1 and ∂_2 need not be orthogonal to each other. Also note that by replacing $\tau \rightarrow \alpha\tau$, α a constant number, in Eq. (17), the normalization condition (28) will change to $h_{00} + \alpha^2\bar{h} \approx 0$.

Using the quasi-orthonormal conditions (27) and (28), the Lagrangian density becomes

$$\mathcal{L}_{NG} \approx -T\bar{h} \approx Th_{00} \approx \frac{T}{2} (h_{00} - \bar{h}).$$

The effective action thus becomes

$$S^{\text{eff}} = \frac{T}{2} \int_{\Sigma} d^3\sigma [h_{00} - h_{11}h_{22} + (h_{12})^2], \quad (29)$$

²One can generalize this gauge fixing condition (24) for higher-dimensional hyper-membranes. Any n -dimensional divergenceless vector field A^a , subjected to the boundary condition $A^a|_{\partial\Sigma} = 0$ (just like $\lambda \cdot \mathcal{P}^a$ in (22) and (23)) can be expressed as $A^a = \varepsilon^{abc_1\dots c_{n-2}}\partial_b B_{c_1\dots c_{n-2}}$, where $B_{c_1\dots c_{n-2}}$ are the components of an $(n-2)$ -form. Like the Kalb-Ramond gauge fields, these B 's have a hierarchy of "gauge symmetries" given by $B \rightarrow B' = B + dB_{(n-3)}$, $B_{(n-3)} \rightarrow B'_{(n-3)} = B_{(n-3)} + dB_{(n-4)}, \dots$, so on and so forth, where $B_{(p)}$ is a p -form. One can therefore easily see that the demand $A^a = 0$ entails $(n-1)$ additional constraints as there are $(n-1)$ independent components of $B_{(n-2)}$. With two gauge fixing conditions of type (17) and (18), this gives rise to $(n+1)$ number of independent constraints, which exactly matches with the number of first-class constraints of the type (7) and (8) of the theory. For the special case of $n = 2$, $A^a = \varepsilon^{ab}\partial_b B$, where B is now a pseudo-scalar. Clearly the demand $A^a = 0$ is equivalent to the gauge fixing condition (24). For the case $n = 3$, $A^a = \varepsilon^{abc}\partial_b B_c$ so that 3-vector is expressed as a curl of another 3-vector, in a standard manner, having only two transverse degrees of freedom; the longitudinal one having been eliminated through the above mentioned gauge transformation.

which gives the equation of motion:

$$\partial_0 \partial_0 X_\mu + \partial_1 (h_{12} \partial_2 X_\mu - h_{22} \partial_1 X_\mu) + \partial_2 (h_{12} \partial_1 X_\mu - h_{11} \partial_2 X_\mu) = 0. \quad (30)$$

Note that the quasi-orthonormal conditions (27) and (28) do not correspond to any gauge conditions themselves as they contain time derivatives. Actually they follow as a consequence of the conditions (17), (18) and (24) which are to be regarded as gauge fixing conditions. These gauge conditions, when imposed, render the first-class constraints (7) and (8) of the theory into second-class as can be seen from their non-vanishing Poisson-bracket structure. Therefore, NG formalism requires the evaluation of Dirac brackets where these constraints are implemented strongly. As we shall see subsequently, in the Polyakov formulation the constraints (7) and (8) are not rendered into second-class and we can avoid the detailed calculation of Dirac brackets.

It is possible to draw a parallel between the quasi-orthonormal gauge discussed here and the usual orthonormal gauge in NG string, which is the analogue of the conformal gauge in the Polyakov string. In the latter case the equations of motion linearize reducing to the D'Alembert equations. This is possible because the gauge choice induces a net of coordinates that form a locally orthonormal system [15]. For the membrane, the invariances are insufficient to make such a choice and the best that we could do was to provide a quasi-orthonormal system. It is however amusing to note that if we forced an orthonormal choice, so that $h_{0a} \approx 0$ is supplemented with $h_{12} \approx 0$ and $h_{11} = h_{22} \approx 1$, then the equation of motion (30) indeed simplifies to the D'Alembert equation. This provides an alternative way of looking at the quasi-orthonormality.

If we do not impose quasi-orthonormality, it is highly non-trivial, if not totally impossible, to express the boundary conditions (4) in terms of phase-space variables because the canonical momentum $\Pi_\mu = \mathcal{P}_\mu^0$ (5), which can be re-expressed as

$$\Pi_\mu = \frac{T\bar{h}}{\sqrt{-h}} \left(\eta_{\mu\nu} - \partial_a X_\mu \bar{h}^{ab} \partial_b X_\nu \right) \partial_0 X^\nu$$

involves a projection operator given by the expression within the parentheses in the above equation. The velocity terms appear both in the right of the projection operator and in $\sqrt{-h}$ appearing in the denominator. This makes the inversion of the above equation to write the velocities in terms of momenta highly non-trivial. Nevertheless, all this simplifies drastically in the quasi-orthonormal gauge to enable us to simplify the above expression to

$$\Pi_\mu = T \partial_0 X_\mu \quad (31)$$

so that the boundary condition (4) is now expressible in terms of phase-space variables as

$$(h_{22} \partial_1 X_\mu - h_{12} \partial_2 X_\mu) \Pi^2 \Big|_{\sigma^1=0, \pi} = 0.$$

Finally we notice that the parameters U^0_0 and V^{a0}_0 given by Eq. (13) simplify in this gauge to

$$U^0_0 = \frac{1}{2T}, \quad V^{a0}_0 = 0 \quad (32)$$

while U^0_a and V^{b0}_a given by Eq. (14) remain unchanged. Now the generators of τ and σ^a translations (12) become

$$H_T = \frac{1}{2T} \int d^2 \sigma \psi, \quad H_a = \int d^2 \sigma \phi_a. \quad (33)$$

It is straightforward to reproduce the action (29) by performing an inverse Legendre transformation. Computing the Poisson bracket of $X_\mu(\tau, \vec{\sigma})$ with the above H_T , the Hamilton's equation $\partial_0 X_\mu = \{X_\mu, H_T\}$ gives Eq. (31), the definition of momenta in this gauge. Then,

$$S^{\text{eff}} = \int_{\Sigma} d^3\sigma \Pi_\mu \partial_0 X^\mu - \int d\tau H_T$$

just yields (29). The other equation, $\partial_0 \Pi_\mu = \{\Pi_\mu, H_T\}$, reproduces Eq. (30), which is the Euler-Lagrange equation following from the effective action (29).

Notice that the values of U^0_0 and V^{a0}_0 are gauge dependent. The particular values given by Eq. (32) correspond to our quasi-orthonormal gauge. Had we chosen a different gauge, we would have obtained different values for these parameters. On the contrary, the parameters U^0_a and V^{b0}_a are gauge independent. This is consistent with the symmetries of the problem. There are three reparametrization invariances, so that three parameters among these U 's and V 's must be gauge dependent, manifesting these symmetries. Since the reparametrization invariances govern the time evolution of the system, the gauge-dependent parameters are given by U^0_0 and V^{a0}_0 , while the others are gauge independent.

3 The Free Polyakov Membrane

The Polyakov action for the bosonic membrane is [13]

$$S_P = -\frac{T}{2} \int_{\Sigma} d^3\sigma \sqrt{-g} \left(g^{ij} \partial_i X^\mu \partial_j X_\mu - 1 \right), \quad (34)$$

where an auxiliary metric g_{ij} on the membrane world-volume has been introduced and will be given the status of an independent field variable in the enlarged configuration space. The final term (-1) inside the parentheses does not appear in the analogous string theory action. A consistent set of equations can be obtained only by taking the ‘‘cosmological’’ constant to be (-1) . Indeed, the equations of motion following from the action (34) but with arbitrary cosmological constant Λ are

$$\partial_i \left(\sqrt{-g} g^{ij} \partial_j X^\mu \right) = 0, \quad (35)$$

$$h_{ij} = \frac{1}{2} g_{ij} \left(g^{kl} h_{kl} + \Lambda \right) \quad (36)$$

while the boundary conditions are

$$\partial^a X^\mu|_{\partial\Sigma} = 0. \quad (37)$$

Eq. (36) can now be satisfied if and only if we identify g_{ij} with h_{ij} :

$$g_{ij} = h_{ij} \equiv \partial_i X^\mu \partial_j X_\mu \quad (38)$$

for the case $\Lambda = -1$ so that the action (34) reduces to the NG action (1). The canonical momenta corresponding to the fields X^μ and g_{ij} are

$$\Pi_\mu = \frac{\partial \mathcal{L}}{\partial \dot{X}^\mu} = -T \sqrt{-g} \partial^0 X_\mu, \quad (39)$$

$$\pi^{ij} = \frac{\partial \mathcal{L}}{\partial \dot{g}_{ij}} = 0. \quad (40)$$

Clearly, $\pi^{ij} \approx 0$ represent primary constraints of the theory. The canonical Hamiltonian density is

$$\begin{aligned}\mathcal{H}_C &= \Pi_\mu \partial_0 X^\mu - \mathcal{L} \\ &= \frac{\sqrt{-g}}{2T\bar{g}} \Pi^2 - \frac{gg^{0a}}{\bar{g}} \Pi_\mu \partial_a X^\mu + \frac{T\sqrt{-g}}{2\bar{g}} (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12} - \bar{g}).\end{aligned}\quad (41)$$

Therefore, the total Hamiltonian is written as

$$H_T = \int d^2\sigma \left(\mathcal{H}_C + \lambda_{ij} \pi^{ij} \right), \quad (42)$$

where λ_{ij} are arbitrary Lagrange multipliers. Conserving the constraint $\pi^{00} \approx 0$ with time:

$$\dot{\pi}^{00} = \{ \pi^{00}, H_T \} \approx 0,$$

we get

$$\Omega_1 \equiv \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12} - \bar{g}) \approx 0. \quad (43)$$

Similarly, conserving other primary constraints with time, we get

$$\begin{aligned}\Omega_2 &\equiv \frac{\sqrt{-g}}{4T\bar{g}^2} (2g_{22} - \bar{g}g^{11}) \left\{ \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12}) \right\} \\ &\quad - \frac{gg_{22}}{\bar{g}^2} g^{0a} \Pi_\mu \partial_a X^\mu - \frac{g_{02}}{\bar{g}} \Pi_\mu \partial_2 X^\mu - \frac{T\sqrt{-g}}{4\bar{g}} (2h_{22} - \bar{g}g^{11}) \approx 0,\end{aligned}\quad (44)$$

$$\begin{aligned}\Omega_3 &\equiv \frac{\sqrt{-g}}{4T\bar{g}^2} (2g_{11} - \bar{g}g^{22}) \left\{ \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12}) \right\} \\ &\quad - \frac{gg_{11}}{\bar{g}^2} g^{0a} \Pi_\mu \partial_a X^\mu - \frac{g_{01}}{\bar{g}} \Pi_\mu \partial_1 X^\mu - \frac{T\sqrt{-g}}{4\bar{g}} (2h_{11} - \bar{g}g^{22}) \approx 0,\end{aligned}\quad (45)$$

$$\begin{aligned}\Omega_4 &\equiv -\frac{\sqrt{-g}}{2T\bar{g}^2} (2g_{12} + \bar{g}g^{12}) \left\{ \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12}) \right\} \\ &\quad + \frac{2gg_{12}}{\bar{g}^2} g^{0a} \Pi_\mu \partial_a X^\mu + \frac{g_{02}}{\bar{g}} \Pi_\mu \partial_1 X^\mu + \frac{g_{01}}{\bar{g}} \Pi_\mu \partial_2 X^\mu + \frac{T\sqrt{-g}}{2\bar{g}} (2h_{12} + \bar{g}g^{12}) \approx 0\end{aligned}\quad (46)$$

$$\begin{aligned}\Omega_5 &\equiv -\frac{\sqrt{-g}g^{01}}{2T} \left\{ \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12} - \bar{g}) \right\} \\ &\quad - g_{22} \Pi_\mu \partial_1 X^\mu + g_{12} \Pi_\mu \partial_2 X^\mu \approx 0,\end{aligned}\quad (47)$$

$$\begin{aligned}\Omega_6 &\equiv -\frac{\sqrt{-g}g^{02}}{2T} \left\{ \Pi^2 + T^2 (g_{22}h_{11} + g_{11}h_{22} - 2g_{12}h_{12} - \bar{g}) \right\} \\ &\quad - g_{11} \Pi_\mu \partial_2 X^\mu + g_{12} \Pi_\mu \partial_1 X^\mu \approx 0.\end{aligned}\quad (48)$$

The above constraints appear to have a complicated form. Also, their connection with the constraints obtained in the NG formalism, is not particularly transparent. To bring the constraints into a more tractable form and to illuminate this connection, it is desirable to express them by the following combinations:

$$\Omega_1 = \psi - T^2 \bar{\chi} \approx 0, \quad (49)$$

$$\Omega_2 = \frac{\sqrt{-g}}{4T\bar{g}^2} (2g_{22} - \bar{g}g^{11}) \Omega_1 - \frac{gg_{22}}{\bar{g}^2} g^{0a} \phi_a - \frac{g_{02}}{\bar{g}} \phi_2 + \frac{T\sqrt{-g}}{2\bar{g}} \chi_{22} \approx 0, \quad (50)$$

$$\Omega_3 = \frac{\sqrt{-g}}{4T\bar{g}^2} (2g_{11} - \bar{g}g^{22}) \Omega_1 - \frac{gg_{11}}{\bar{g}^2} g^{0a} \phi_a - \frac{g_{01}}{\bar{g}} \phi_1 + \frac{T\sqrt{-g}}{2\bar{g}} \chi_{11} \approx 0, \quad (51)$$

$$\Omega_4 = -\frac{\sqrt{-g}}{2T\bar{g}^2} (2g_{12} + \bar{g}g^{12}) \Omega_1 + \frac{2gg_{12}}{\bar{g}^2} g^{0a} \phi_a + \frac{g_{02}}{\bar{g}} \phi_1 + \frac{g_{01}}{\bar{g}} \phi_2 - \frac{T\sqrt{-g}}{\bar{g}} \chi_{12} \approx 0, \quad (52)$$

$$\Omega_5 = -\frac{\sqrt{-g}g^{01}}{2T} \Omega_1 - g_{22}\phi_1 + g_{12}\phi_2 \approx 0, \quad (53)$$

$$\Omega_6 = -\frac{\sqrt{-g}g^{02}}{2T} \Omega_1 - g_{11}\phi_2 + g_{12}\phi_1 \approx 0, \quad (54)$$

where

$$\psi \equiv \Pi^2 + T^2 \bar{h} \approx 0, \quad (55)$$

$$\phi_a \equiv \Pi_\mu \partial_a X^\mu \approx 0, \quad (56)$$

$$\chi_{ab} \equiv g_{ab} - h_{ab} \approx 0 \quad (57)$$

and $\bar{\chi} = \chi_{11}\chi_{22} - (\chi_{12})^2$. As all the constraints Ω 's appearing in Eqs. (49–54) are combinations of ψ , ϕ_a and χ_{ab} in Eqs. (55–57), we can treat these ψ , ϕ_a and χ_{ab} as an alternative set of secondary constraints. These constraints along with the primary constraints $\pi^{ij} \approx 0$ (40) constitute the complete set of constraints of the theory. This is because the canonical Hamiltonian density (41) can be expressed as a combination of constraints in the following manner:

$$\mathcal{H}_C = \frac{\sqrt{-g}}{2T\bar{g}} \psi - \frac{gg^{0a}}{\bar{g}} \phi_a - \frac{T\sqrt{-g}}{2\bar{g}} \bar{\chi} \approx 0 \quad (58)$$

and the non-vanishing Poisson brackets between the constraints of the theory are

$$\begin{aligned} \{\psi(\tau, \vec{\sigma}), \chi_{ab}(\tau, \vec{\sigma}')\} &\approx 2 (\partial_a \Pi_\mu \partial_b X^\mu + \partial_b \Pi_\mu \partial_a X^\mu) \delta(\vec{\sigma} - \vec{\sigma}'), \\ \{\phi_a(\tau, \vec{\sigma}), \chi_{bc}(\tau, \vec{\sigma}')\} &= h_{ab}(\tau, \vec{\sigma}') \partial'_c \delta(\vec{\sigma} - \vec{\sigma}') + h_{ac}(\tau, \vec{\sigma}') \partial'_b \delta(\vec{\sigma} - \vec{\sigma}') \\ &\quad + (\partial_b X^\mu \partial_c \partial_a X_\mu + \partial_c X^\mu \partial_b \partial_a X_\mu) \delta(\vec{\sigma} - \vec{\sigma}'), \\ \{\pi^{ab}(\tau, \vec{\sigma}), \chi_{cd}(\tau, \vec{\sigma}')\} &= -\frac{1}{2} (\delta_c^a \delta_d^b + \delta_d^a \delta_c^b) \delta(\vec{\sigma} - \vec{\sigma}'), \end{aligned} \quad (59)$$

while the weakly vanishing brackets are the same as given by (9). As far as the rest of the brackets are concerned, it is trivial to see that they vanish strongly. Thus, as it appears, none of the constraints except π^{0i} in the set is first-class. But we have not yet extracted the maximal number of first-class constraints from the set (40, 55–57) by constructing appropriate linear combinations of the constraints. However, it is highly non-trivial to find such a linear combination in the present case as one can see from the complicated structure of the Poisson brackets given above in (59). Nevertheless, one can bypass such an elaborate procedure to extract the first-class constraints from the given set by noting that the complete set of constraints can be split into two sectors. In one sector we retain ψ , ϕ_a and π^{0i} , which are first-class among themselves, while the other sector contains the canonically conjugate pairs χ_{ab} and π^{ab} . This allows an iterative computation of the Dirac brackets [16]; namely, it is possible to eliminate this set completely by calculating the Dirac brackets within this sector. The brackets of the other constraints are now computed with respect to these Dirac brackets. Obviously ψ , ϕ_a will have vanishing brackets with π^{ab} , χ_{cd} . Moreover, the original first-class algebra among ψ and ϕ_a will be retained. This follows from the fact that the Dirac constraint matrix involving π^{ab} and χ_{cd} has entries only in

the off-diagonal pieces, while ψ and ϕ_a have non-vanishing contributions coming just from the bracket with one of them; i.e. χ_{cd} (see (59)). The Dirac brackets of ψ and ϕ_a are thus identical to their Poisson brackets, satisfying the same algebra as in the NG case.

We are therefore left with the first-class constraints $\psi \approx 0$, $\phi_a \approx 0$ and $\pi^{0i} \approx 0$. At this stage, we note that the constraints $\pi^{0i} \approx 0$ are analogous to $\pi^0 \approx 0$ in free Maxwell theory, where π^0 is canonically conjugate to A_0 . Consequently, the time evolution of g_{0i} is arbitrary as follows from the Hamiltonian (42). Therefore, we can set

$$g_{0a} = 0, \quad g_{00} = -\bar{h}, \quad (60)$$

as new gauge fixing conditions.³ With that (g_{0a}, π^{0a}) and (g_{00}, π^{00}) are discarded from the phase-space. This is again analogous to the arbitrary time evolution of A_0 in Maxwell theory, where we can set $A_0 = 0$ as a gauge fixing condition and discard the pair (A_0, π^0) from the phase-space altogether.

These gauge fixing conditions (60) are the counterpart of the quasi-orthonormal conditions (27) and (28) in the NG case. However, unlike the NG case, these second-class constraints (60) do not render the residual first-class constraints of the theory, viz. $\psi \approx 0$ and $\phi_a \approx 0$ into second-class constraints. Therefore, they represent partial gauge fixing conditions. This stems from the fact that g_{0i} were still regarded as independent field variables in the configuration space whereas g_{ab} have already been strongly identified with h_{ab} (57). We therefore note that the calculation of the Dirac brackets is not necessary in Polyakov formulation. This motivates us to study the noncommutativity vis-à-vis the modified brackets $\{X^\mu, X^\nu\}$ in the simpler Polyakov version. For that we shall first consider the free theory in the next section.

Before we conclude this section, let us make some pertinent observations about the structure of the symmetric form of energy-momentum tensor, which is obtained by functionally differentiating the action with respect to the metric. The various components of this tensor are given by:

$$\begin{aligned} T_{00} = & \frac{g_{00}}{2T\bar{g}}\psi + \frac{2g\sqrt{-g}}{\bar{g}^2}g^{0a}\phi_a + \left(\frac{1}{g^{00}} - g_{00}\right)\frac{\Pi^2}{T\bar{g}} \\ & - \frac{Tg^2}{\bar{g}^2} \left[(g^{01})^2 h_{11} + (g^{02})^2 h_{22} + 2g^{01}g^{02}h_{12} \right] - \frac{Tg_{00}}{2\bar{g}}\bar{\chi}, \end{aligned} \quad (61)$$

$$T_{01} = -\frac{g_{01}}{2T\bar{g}}\psi - \frac{\sqrt{-g}}{\bar{g}}\phi_1 + \frac{T}{\bar{g}} \left[(g_{02}h_{11} + g_{01}h_{12})\chi_{12} - g_{02}h_{12}\chi_{11} - g_{01}h_{11}\chi_{22} - \frac{g_{01}}{2}\bar{\chi} \right] \quad (62)$$

$$T_{02} = -\frac{g_{02}}{2T\bar{g}}\psi - \frac{\sqrt{-g}}{\bar{g}}\phi_2 + \frac{T}{\bar{g}} \left[(g_{01}h_{22} + g_{02}h_{12})\chi_{12} - g_{01}h_{12}\chi_{22} - g_{02}h_{22}\chi_{11} - \frac{g_{02}}{2}\bar{\chi} \right] \quad (63)$$

$$T_{ab} = -\frac{g_{ab}}{2T\bar{g}}\psi + T\chi_{ab} + \frac{Tg_{ab}}{2\bar{g}}\bar{\chi} - \frac{Tg_{ab}}{\bar{g}}(g_{22}\chi_{11} + g_{11}\chi_{22} - 2g_{12}\chi_{12}). \quad (64)$$

Note that unlike the case of string [11], the component T_{00} cannot be written in terms of constraints of the theory. However, the other components can be expressed in terms of these constraints, of which χ_{ab} are second-class and have already been put strongly to zero by using Dirac brackets, so that the form of T_{0a} and T_{ab} simplifies to

$$T_{0a} = -\frac{g_{0a}}{2T\bar{g}}\psi - \frac{\sqrt{-g}}{\bar{g}}\phi_a,$$

³We cannot set $g_{00} = 0$ as it will make the metric singular. We therefore set $g_{00} = -\bar{h}$ to make it match with the corresponding condition (28) in NG case.

$$T_{ab} = -\frac{g_{ab}}{2T\bar{g}}\psi.$$

However, for T_{00} we have to make use of the gauge conditions (60), which hold strongly as was discussed earlier, to enable us to write

$$T_{00} = -\frac{1}{2T}\psi.$$

Let us now compare it with NG case. First we notice that θ^i_j appearing in Eq. (11) is not a tensor itself but it is a tensor density. The corresponding tensor is $\frac{1}{\sqrt{-g}}\theta^i_j$. In quasi-orthonormal gauge, we have

$$\sqrt{-g}T^0_0 = \theta^0_0 = \frac{1}{2T}\psi,$$

which reproduces the canonical Hamiltonian density (58) in this gauge. Also, in this gauge, we have

$$\sqrt{-g}T^0_a = \phi_a,$$

which matches with θ^0_a in quasi-orthonormal gauge. This also provides a direct generalization of the string case [11]. Although, unlike the string case, the Weyl symmetry is absent in the membrane case, we still have a vanishing trace, albeit weakly, of the energy-momentum tensor:

$$T^i_i = -\frac{1}{2T\bar{h}}\psi \approx 0.$$

4 The Brackets for a Free Theory

Here we consider a cylindrical topology for the membrane which is taken to be periodic along σ^2 -direction, i.e. $\sigma^2 \in [0, 2\pi)$ and $\sigma^1 \in [0, \pi]$. Following the example of string case [11], we write down the first version of the brackets as:

$$\{X^\mu(\tau, \vec{\sigma}), \Pi_\nu(\tau, \vec{\sigma}')\} = \delta^\mu_\nu \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2 - \sigma'^2), \quad (65)$$

and the other brackets vanishing.⁴ Here

$$\delta_P(\sigma - \sigma') = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} e^{in(\sigma - \sigma')} \quad (66)$$

is the periodic delta function of period 2π which satisfies

$$\int_{-\pi}^{+\pi} d\sigma' \delta_P(\sigma - \sigma') f(\sigma') = f(\sigma) \quad (67)$$

for any periodic function $f(\sigma) = f(\sigma + 2\pi)$ defined in the interval $[-\pi, +\pi]$; and if, in addition, $f(\sigma)$ is taken to be an even function in the interval $[-\pi, +\pi]$, then the above integral (67) reduces to

$$\int_0^\pi d\sigma' \Delta_+(\sigma, \sigma') f(\sigma') = f(\sigma), \quad (68)$$

⁴Note that the $\{X^\mu, \Pi_\nu\}$ brackets are not affected as we implemented the second-class constraints and the gauge fixing conditions strongly in the preceding section. They are the only surviving phase-space variables as g_{ij} have lost their independent status.

where

$$\begin{aligned}\Delta_+(\sigma, \sigma') &= \delta_P(\sigma - \sigma') + \delta_P(\sigma + \sigma') \\ &= \frac{1}{\pi} + \frac{1}{\pi} \sum_{n \neq 0} \cos(n\sigma) \cos(n\sigma').\end{aligned}\tag{69}$$

This structure of the brackets is, however, consistent only with Neumann boundary conditions along σ^1 -direction. On the other hand, we have a mixed boundary condition (37) which can be expressed in terms of phase-space variables as

$$\left[g_{22} T \partial_1 X^\mu + \sqrt{-g} g^{01} \Pi^\mu - g_{12} T \partial_2 X^\mu \right]_{\sigma^1=0, \pi} = 0.\tag{70}$$

We notice that in NG formulation it was necessary to fix gauge in order to express the boundary condition in terms of phase-space variables. However, this is not the case with Polyakov formulation since g_{ij} are taken to be independent fields. Using the strongly valid equations (57) and the gauge fixing conditions (60), this simplifies further to

$$[\partial_2 X^\nu \partial_2 X_\nu \partial_1 X^\mu - \partial_1 X^\nu \partial_2 X_\nu \partial_2 X^\mu]_{\sigma^1=0, \pi} = 0.\tag{71}$$

Although we are using the gauge (60), the non-trivial gauge generating first-class constraints (55) and (56) will be retained in the gauge-independent analysis both here and in the interacting case. As we see, the above boundary condition is non-trivial in nature and involves both the ∂_1 and ∂_2 derivatives. But, since the coordinates and momenta are not related at the boundary, we do not require to postulate a non-vanishing $\{X^\mu, X^\nu\}$ bracket as in the case of free string in conformal gauge [11]. Therefore, the free membrane theory, like its string counterpart, does not exhibit noncommutativity in the boundary coordinates.

5 The Low-Energy Limit

In this section, we would like to see how the results in the free membrane theory go over to those of free string theory in the limit of small radius for the cylindrical membrane.

The cylindrical membrane is usually taken to propagate in an 11-dimensional compactified target space $\mathbf{R}^{9-p} \times M^p \times S^1 \times I$, where M^p is a p -dimensional flat Minkowski space-time and I is an interval with finite length. There exist at the boundaries of I two p -branes on which an open membrane can end. And the topology of the p -branes is given by $M^p \times S^1$. Also, the cylindrical membrane is assumed to wrap around this S^1 . The radius of this circle is supposed to be very small so that in the low-energy limit the target space effectively goes over to 10-dimensional $\mathbf{R}^{9-p} \times M^p \times I$ and the cylindrical membrane goes over to the open string.

At this stage, we choose further gauge fixing conditions:

$$X^0 = \tau, \quad X^2 = \sigma^2 R,\tag{72}$$

where we have introduced R to indicate the radius of the cylindrical membrane and X^2 represents the compact dimension S^1 .⁵ Before choosing the gauge conditions (72), the τ

⁵In [6], another gauge fixing condition $X^1 = \sigma^1$, (π being the length of the cylindrical membrane) has been used. But we notice that imposition of this gauge fixing condition would be inconsistent with the boundary condition (71) since, for $\mu = 1$, it yields a topology changing condition (cylinder $\rightarrow$ sphere), $R^2|_{\sigma^1=0, \pi} = 0$, which is clearly unacceptable. Therefore, the choice (72) does not allow us to choose $X^1 = \sigma^1$ as well, which is not needed either for our purpose.

and σ^a translations were generated by the constraints $\frac{1}{2T}\psi$ and ϕ_a respectively, just as in the NG case (33). Now we have

$$\begin{aligned}\left\{\psi(\tau, \vec{\sigma}), X^0(\tau, \vec{\sigma}') - \tau\right\} &= -2\Pi^0(\tau, \vec{\sigma}) \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2, \sigma'^2), \\ \left\{\phi_2(\tau, \vec{\sigma}), X^2(\tau, \vec{\sigma}') - \sigma'^2 R\right\} &= -\partial_2 X^2(\tau, \vec{\sigma}) \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2, \sigma'^2) \\ &\approx -R \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2, \sigma'^2),\end{aligned}$$

whereas

$$\begin{aligned}\left\{\phi_1(\tau, \vec{\sigma}), X^0(\tau, \vec{\sigma}') - \tau\right\} &= -\partial_1 X^0(\tau, \vec{\sigma}) \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2, \sigma'^2) \approx 0, \\ \left\{\phi_1(\tau, \vec{\sigma}), X^2(\tau, \vec{\sigma}') - \sigma'^2 R\right\} &= -\partial_1 X^2(\tau, \vec{\sigma}) \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2, \sigma'^2) \approx 0.\end{aligned}$$

Thus, the (partial) gauge fixing conditions (72) take care of the world-volume diffeomorphism generated by ψ and ϕ_2 in the sense that these constraints are rendered into second-class while the diffeomorphism generated by ϕ_1 is still there.

Coming back to the low-energy limit, we would like to show that the σ^2 dependence of all the fields except X^2 itself drops out effectively in the gauge (72). To motivate it, let us consider the case of a free massless scalar field defined on a space with one compact dimension of ignorable size. Let the space be $M^p \times S^1$, where M^p is a p -dimensional Minkowski space-time taken to be flat for simplicity and S^1 is a circle of radius R which is very small. We take $\theta \in [0, 2\pi)$ to be the angle coordinate corresponding to this circle so that the metric is given by $ds^2 = \eta_{\mu\nu} dx^\mu dx^\nu = \eta_{\mu'\nu'} dx^{\mu'} dx^{\nu'} + R^2 d\theta^2$ with μ, ν ranging from 0 to p and μ', ν' from 0 to $(p-1)$. The action is

$$S = -\frac{1}{2} \int d^p x d\theta \partial_\mu \phi \partial^\mu \phi.$$

Separating the index corresponding to the compact dimension, we rewrite it as

$$S = -\frac{1}{2} \int d^p x d\theta \left(\partial_{\mu'} \phi \partial^{\mu'} \phi + \frac{1}{R^2} \partial_\theta \phi \partial_\theta \phi \right).$$

Substituting the Fourier expansion

$$\phi(x, \theta) = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} \phi_{(n)}(x) e^{in\theta}, \quad \phi_{(-n)} = \phi_{(n)}^*$$

in the action and integrating out the compact dimension, we get

$$S \rightarrow S' = -\frac{1}{2} \int d^p x \sum_{n \in \mathbb{Z}} \left(\partial_{\mu'} \phi_{(n)} \partial^{\mu'} \phi_{(n)}^* + \frac{n^2}{R^2} \phi_{(n)} \phi_{(n)}^* \right).$$

Thus the Fourier coefficients represent a whole tower of effective massive complex scalar fields of mass $\sim \frac{n}{R}$ in a lower-dimensional non-compact space-time. These masses are usually of the Planck order if R is of the order of Planck length and are therefore ignored in the low-energy regime. Equivalently, one ignores the θ dependence of the field ϕ . This can also be understood from physical considerations. In the low-energy limit, the associated wavelengths are very large as compared to R so that variation of the field along the circle is ignorable.

Now the membrane goes over to string in the low-energy regime when the circle S^1 effectively disappears in the limit $R \rightarrow 0$. So the field theory living in the membrane world-volume is expected to correspond to the field theory living on string world-sheet. To verify this, let us substitute the Fourier expansion of the world-volume fields $X^\mu(\tau, \sigma^1, \sigma^2)$ around σ^2 :

$$X^\mu(\tau, \sigma^1, \sigma^2) = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} X_{(n)}^\mu(\tau, \sigma^1) e^{in\sigma^2}, \quad X_{(-n)}^\mu = X_{(n)}^{\mu *} \quad (73)$$

in the Poisson bracket (65) to find that the Fourier coefficients $X_{(n)}^\mu(\tau, \sigma^1)$ satisfy

$$\{X_{(n)}^\mu(\tau, \sigma^1), \Pi_\nu^{(m)}(\tau, \sigma'^1)\} = \delta_\nu^\mu \delta_n^{-m} \Delta_+(\sigma^1, \sigma'^1). \quad (74)$$

As in the case of free scalar field discussed above, the Fourier coefficients $X_{(0)}^\mu(\tau, \sigma^1)$ will represent the effective (real) fields in the string world-sheet satisfying

$$\{X_{(0)}^\mu(\tau, \sigma^1), \Pi_\nu^{(0)}(\tau, \sigma'^1)\} = \delta_\nu^\mu \Delta_+(\sigma^1, \sigma'^1), \quad (75)$$

which reproduces the Poisson bracket for string (see appendix). The sub/superscript (0) will be dropped now onwards for convenience. Using $\partial_2 X^\mu = R\delta_2^\mu$, the boundary condition (71) gives

$$\partial_1 X^\mu|_{\sigma^1=0, \pi} = 0 \quad (76)$$

so that we recover the boundary condition for free string in conformal gauge (see appendix).⁶

Now we would like to show how the gauge fixed world-volume membrane metric (60) reduces to the world-sheet string metric in conformal gauge. For that first note that the components of the metric tensor in a matrix form can be written as

$$\{g_{ij}\} = \begin{pmatrix} g_{00} & g_{01} & g_{02} \\ g_{01} & g_{11} & g_{12} \\ g_{02} & g_{12} & g_{22} \end{pmatrix} = \begin{pmatrix} g_{00} & 0 & 0 \\ 0 & h_{11} & 0 \\ 0 & 0 & R^2 \end{pmatrix},$$

where we have made use of the first gauge fixing condition in (60) and by now the strongly valid equations (57). Clearly, this matrix becomes singular in the limit $R \rightarrow 0$ taken in a proper mathematical sense. It must therefore correspond to a two-dimensional surface embedded in three-dimensional world-volume. The metric corresponding to it can be easily obtained by chopping off the last row and last column in the above three-dimensional metric to get $\begin{pmatrix} g_{00} & 0 \\ 0 & h_{11} \end{pmatrix}$. Now, we make use of the second gauge fixing condition in (60) to replace g_{00} by $(-\bar{h})$. However, this $\bar{h}$ can be simplified further using the gauge (72) to get $R^2 h_{11}$ so that the above 2×2 matrix becomes $h_{11} \begin{pmatrix} -R^2 & 0 \\ 0 & 1 \end{pmatrix}$ and the diagonal elements get identified up to a scale factor. It can now be put in the standard form, $\text{diag}(-1, 1)$, upto an overall Weyl factor, by replacing the second condition in (60) by $g_{00} = -\alpha^2 \bar{h}$ and choosing α suitably. We also notice that using $h_{0a} = 0$, the NG action for the membrane becomes

$$S_{NG} = -T \int d^3\sigma \sqrt{-h_{00} \bar{h}},$$

⁶Actually, we do not get (76) directly, rather it is accompanied by a pre-factor R^2 . However, this equation is not satisfied trivially if $R \rightarrow 0$, as this limit should not be taken literally in a mathematical sense. This just means that R should be taken to have a very small non-zero value and presumably should be of the order of Planck length, as we have mentioned earlier.

which using the gauge conditions (72) and integrating out σ^2 , reduces to the NG action for string in orthonormal gauge:

$$S_{NG} \rightarrow S'_{NG} = -2\pi RT \int d^2\sigma \sqrt{-h_{00}h_{11}}.$$

This also shows that the string tension is $\sim TR$ if the original membrane tension is given by T . Actually one takes the limit $R \rightarrow 0$ together with the membrane tension $T \rightarrow \infty$ in such a way that their product (TR) is finite. Such a limit was earlier discussed, from other considerations, in [17].

6 The Interacting Membrane

The Polyakov action for a membrane moving in the presence of a constant antisymmetric background field $A_{\mu\nu\rho}$ is

$$S_P = -\frac{T}{2} \int_{\Sigma} d^3\sigma \left[\sqrt{-g} \left(g^{ij} \partial_i X^\mu \partial_j X_\mu - 1 \right) + \frac{e}{3} \varepsilon^{ijk} \partial_i X^\mu \partial_j X^\nu \partial_k X^\rho A_{\mu\nu\rho} \right], \quad (77)$$

where we have introduced a coupling constant e .⁷ The equations of motion are

$$\partial_i \left(\sqrt{-g} g^{ij} \partial_j X_\mu + \frac{e}{2} \varepsilon^{ijk} \partial_j X^\nu \partial_k X^\rho A_{\mu\nu\rho} \right) = 0, \quad (78)$$

$$g_{ij} = h_{ij} \equiv \partial_i X^\mu \partial_j X_\mu. \quad (79)$$

Note that the second equation does not change from the free case ($e = 0$) despite the presence of interaction term as this term is topological in nature and does not involve the metric g_{ij} . The canonical momenta are

$$\Pi_\mu = \frac{\partial \mathcal{L}}{\partial \dot{X}^\mu} = -T \left(\sqrt{-g} \partial^0 X_\mu + \frac{e}{2} \varepsilon^{ab} \partial_a X^\nu \partial_b X^\rho A_{\mu\nu\rho} \right), \quad (80)$$

$$\pi^{ij} = \frac{\partial \mathcal{L}}{\partial \dot{g}_{ij}} = 0. \quad (81)$$

For convenience, we define

$$\tilde{\Pi}_\mu \equiv \Pi_\mu + \frac{eT}{2} \varepsilon^{ab} \partial_a X^\nu \partial_b X^\rho A_{\mu\nu\rho} = -T \sqrt{-g} \partial^0 X_\mu. \quad (82)$$

Proceeding just as in the free case, the structure of the Hamiltonian density $\mathcal{H}_C$ and the set of constraints is obtained just by replacing $\Pi_\mu \rightarrow \tilde{\Pi}_\mu$, so that we are finally left with the following first-class constraints:

$$\psi \equiv \tilde{\Pi}^2 + T^2 \bar{h} \approx 0, \quad (83)$$

$$\phi_a \equiv \tilde{\Pi}_\mu \partial_a X^\mu \approx 0 \quad (84)$$

⁷As it stands, the interaction term involving the three-form field $A_{\mu\nu\rho}$ in (77) is not gauge invariant under the transformation $A \rightarrow A + d\Lambda$, where Λ is a two-form field. One can, however, make it gauge invariant by adding a surface term $2e \int_{\partial\Sigma} B$, where B is a two-form undergoing the compensating gauge transformation $B \rightarrow B - \Lambda$. But, using Stoke's theorem, this gets combined to a single integral over the world-volume as $\int_{\Sigma} (A + dB)$ so that $(A + dB)$ is gauge invariant as a whole. In the action (77), A is taken to correspond to this gauge invariant quantity by absorbing dB in A .

and, as argued in the free case, we adopt the same gauge fixing conditions (60).

For a cylindrical membrane periodic along σ^2 -direction with $\sigma^1 \in [0, \pi]$, $\sigma^2 \in [0, 2\pi)$, the boundary condition is given by

$$\left[\sqrt{-g} \partial^1 X_\mu + e \partial_2 X^\nu \partial_0 X^\rho A_{\mu\nu\rho} \right]_{\sigma^1=0,\pi} = 0, \quad (85)$$

which when expressed in terms of phase-space variables looks as

$$\left[g_{22} T \partial_1 X_\mu - g_{12} T \partial_2 X_\mu + \sqrt{-g} g^{01} \Pi_\mu + e \left(\Pi^\rho + e T \partial_1 X^\lambda \partial_2 X^\kappa A^\rho_{\lambda\kappa} \right) \partial_2 X^\nu A_{\mu\nu\rho} \right]_{\sigma^1=0,\pi} = 0. \quad (86)$$

As in the free case, here also we use the strongly valid equations (57) and the gauge fixing conditions (60) so that the above boundary condition simplifies to

$$\left[T \partial_2 X^\nu \partial_2 X_\nu \partial_1 X_\mu - T \partial_1 X^\nu \partial_2 X_\nu \partial_2 X_\mu + e \left(\Pi^\rho + e T \partial_1 X^\lambda \partial_2 X^\kappa A^\rho_{\lambda\kappa} \right) \partial_2 X^\nu A_{\mu\nu\rho} \right]_{\sigma^1=0,\pi} = 0. \quad (87)$$

Here we notice that the above boundary condition involves both phase-space coordinates (X^μ, Π_ν) . Using the brackets of the free theory to compute the Poisson bracket of the left-hand side of above equation with $X_\sigma(\tau, \vec{\sigma}')$, we find that it does not vanish. The boundary condition is therefore not compatible with the brackets of the free theory. Thus, we have to postulate a non-vanishing $\{X^\mu, X^\nu\}$ bracket.⁸ For that we make an ansatz:

$$\begin{aligned} \{X_\mu(\tau, \vec{\sigma}), X_\nu(\tau, \vec{\sigma}')\} &= C_{\mu\nu}(\vec{\sigma}, \vec{\sigma}') \\ &= C_{\mu\nu}(\sigma^1, \sigma'^1) \delta_P(\sigma^2 - \sigma'^2) \end{aligned} \quad (88)$$

with

$$C_{\mu\nu}(\sigma^1, \sigma'^1) = -C_{\nu\mu}(\sigma'^1, \sigma^1). \quad (89)$$

and the $\{X^\mu, \Pi_\nu\}$ bracket is taken to be the same as in the free case — Eq. (65). At this stage, we note that the boundary condition (87), if bracketted with $X_\sigma(\tau, \vec{\sigma}')$, yields at the boundary

$$\begin{aligned} &\left[T \partial_2 X^\nu \partial_2 X_\nu \delta_\mu^\lambda - T \partial_2 X^\lambda \partial_2 X_\mu + e^2 T \partial_2 X^\kappa \partial_2 X^\nu A_{\mu\nu\rho} A^{\rho\lambda}_\kappa \right] \partial_1 C_{\lambda\sigma}(\vec{\sigma}, \vec{\sigma}') \\ &+ \left[2T \partial_1 X_\mu \partial_2 X^\kappa - T \partial_1 X^\nu \partial_2 X_\nu \delta_\mu^\kappa - T \partial_2 X_\mu \partial_1 X^\kappa + e \Pi_\rho A_\mu^{\kappa\rho} \right. \\ &\quad \left. + e^2 T \partial_1 X^\lambda \partial_2 X^\nu \left(A_{\mu\nu\rho} A^\rho_{\lambda\kappa} + A_\mu^{\kappa\rho} A^\rho_{\lambda\nu} \right) \right] \partial_2 C_{\kappa\sigma}(\vec{\sigma}, \vec{\sigma}') \\ &= e \partial_2 X^\nu A_{\mu\nu\sigma} \Delta_+(\sigma^1, \sigma'^1) \delta_P(\sigma^2 - \sigma'^2), \end{aligned} \quad (90)$$

which involves both $\partial_1 \mathcal{C}$ and $\partial_2 \mathcal{C}$ and leads to a contradiction if we put $C_{\mu\nu}(\vec{\sigma}, \vec{\sigma}') = 0$. This is another way of seeing that there must be a noncommutativity in the membrane coordinates. However, there is no contradiction with $C_{\mu\nu}(\vec{\sigma}, \vec{\sigma}') = 0$ provided $A_{\mu\nu\rho} = 0$, thereby implying that there is no noncommutativity in the free theory.

Because of the non-linearity in the above equation, it is problematic to find an exact solution. It should however be stressed that the above relation has been derived in a general (gauge-independent) manner. At this point there does not seem to be any compelling reason to choose a particular gauge to simplify this equation further to enable an exact

⁸In the case of free Polyakov string also, the incompatibility of the boundary condition with the basic Poisson brackets forces us to postulate a non-vanishing $\{X^\mu, X^\nu\}$. However, in contrast to the interacting string, this bracket vanishes in a particular gauge — the conformal gauge. (See appendix.)

solution. Nonlinearity would, in all probability, prevent this. This is in contrast to the string case where the analysis naturally leads to a class of light-cone gauges where the corresponding equation was solvable [11]. However, by taking recourse to the low-energy approximation, we show that the results for the string case are recovered. To this end, we substitute the expansion (73) in (88) to get

$$\{X_{(n)}^\mu(\tau, \sigma^1), X_{(m)}^\nu(\tau, \sigma'^1)\} = \delta_{n,-m} C^{\mu\nu}(\sigma^1, \sigma'^1). \quad (91)$$

But again, as in the free case, we retain only the real fields $X_{(0)}^\mu(\tau, \sigma^1) \equiv X^\mu(\tau, \sigma^1)$ when we consider the low-energy regime. Using the gauge fixing conditions (72), the boundary condition (87) reduces to

$$\left[(TR) \partial_1 X_\mu - e \Pi^\rho A_{\mu\rho 2} - e^2 (TR) \partial_1 X_\lambda A^{\rho\lambda}{}_2 A_{\mu\rho 2} \right]_{\sigma^1=0,\pi} = 0. \quad (92)$$

Here, X^μ and Π_ν can be taken to correspond to $X_{(0)}^\mu(\tau, \sigma^1)$ and $\Pi_\nu^{(0)}(\tau, \sigma^1)$ respectively. Thus we recover the boundary condition of the string theory in conformal gauge with the correspondence $TR \leftrightarrow T_S$ and $A_{\mu\nu 2} \leftrightarrow B_{\mu\nu}$, where T_S is the effective (string) tension and $B_{\mu\nu}$ is the 2-form background field appearing in the string theory [11]. Now taking the Poisson bracket of the boundary condition (92) with $X_\sigma(\tau, \sigma^1)$, the low-energy effective real fields, one gets for $\mu \neq 2$ the following differential condition satisfied by $C_{\mu\sigma}$ at the boundary

$$T_S \left(\delta_\mu^\lambda - e^2 A_{\mu\rho 2} A^{\rho\lambda}{}_2 \right) \partial_1 C_{\lambda\sigma}(\sigma^1, \sigma'^1) \Big|_{\sigma^1=0,\pi} = e A_{\sigma\mu 2} \Delta_+(\sigma^1, \sigma'^1) \Big|_{\sigma^1=0,\pi}, \quad (93)$$

which just reproduces the corresponding equation in string theory (see appendix, in particular equation (126)). We therefore obtain the noncommutativity:

$$C_{\mu\nu}(\sigma^1, \sigma'^1) = \frac{1}{2} (NM^{-1})_{(\nu\mu)} [\Theta(\sigma^1, \sigma'^1) - \Theta(\sigma'^1, \sigma^1)] + \frac{1}{2} (NM^{-1})_{[\nu\mu]} [\Theta(\sigma^1, \sigma'^1) + \Theta(\sigma'^1, \sigma^1) - 1], \quad (94)$$

where

$$N_{\nu\sigma} = e A_{\nu\sigma 2}, \quad M^\lambda{}_\mu = T_S \left(\delta_\mu^\lambda - e^2 A_{\mu\rho 2} A^{\rho\lambda}{}_2 \right)$$

with $(NM^{-1})_{(\nu\mu)}$ the symmetric and $(NM^{-1})_{[\nu\mu]}$ the antisymmetric part of $(NM^{-1})_{\nu\mu}$ while

$$\Theta(\sigma^1, \sigma'^1) = \frac{\sigma^1}{\pi} + \frac{1}{\pi} \sum_{n \neq 0} \frac{1}{n} \sin(n\sigma^1) \cos(n\sigma'^1) \quad (95)$$

being the generalized step function which satisfies

$$\partial_1 \Theta(\sigma^1, \sigma'^1) = \Delta_+(\sigma^1, \sigma'^1). \quad (96)$$

It has the property

$$\begin{aligned} \Theta(\sigma^1, \sigma'^1) &= 1 \quad \text{for } \sigma^1 > \sigma'^1, \\ \Theta(\sigma^1, \sigma'^1) &= 0 \quad \text{for } \sigma^1 < \sigma'^1. \end{aligned}$$

7 Concluding Remarks

We have analysed an open membrane, with square and cylindrical topology, ending on p -branes. Both the free case as well as the theory where the membrane is coupled to a background three-form potential were considered.

For the free theory, the world-volume action was taken to be either the NG type or the Polyakov type. For the NG action, a gauge-independent formulation, similar to that adopted in [12] for the string theory, was presented. The reparametrization invariances were manifested by the freedom in the choice of the multipliers enforcing the constraints of the theory. The implications of the boundary conditions in preserving the stability of the free membrane were discussed, highlighting the parallel with the string treatment. A set of quasi-orthonormal gauge fixing conditions was systematically obtained, which simplified the structure of the Hamiltonian.

A constrained analysis of the Polyakov action, contrary to the NG action, led to the presence of second-class constraints. However, by an iterative prescription of computing Dirac brackets, the first-class sector was identified. The Dirac brackets of this sector were identical to the Poisson brackets and exactly matched with the involutive algebra found in the NG case. The analogue of the quasi-orthonormal gauge was also discussed in the Polyakov formulation. It naturally led to the choice of the metric which is used to perform calculations in the light-front variables [13]. Moreover, in this gauge, the energy-momentum tensor was expressed as a combination of the constraints. On the constraint shell, this tensor was seen to have a vanishing trace.

A fundamental difference of the quasi-orthonormal gauge fixing in the two cases was pointed out. In the Polyakov case, gauge fixing entailed certain restrictions on the metric. Since the metric is regarded as an independent field, the gauge fixing does not affect the constraints of the theory which generate the reparametrization invariances. The discussion was thus confined to the Poisson algebra only. A similar gauge fixing in the NG case obviously restricts the target space coordinates. The first-class constraints get converted into second-class ones, thereby necessitating the use of Dirac brackets. Their evaluation is quite complicated due to non-linear terms.

Since Dirac brackets were avoided in the Polyakov formulation, we proceeded to discuss noncommutativity only in this formulation. Also, cylindrical topology of the membrane was considered. We stress that, contrary to standard approaches [4, 5, 6, 7, 8, 9, 10], boundary conditions were not treated as primary constraints of the theory. Our approach was in line with the previous treatment for string theory [11], which has been summarized in the appendix. Thus, noncommutative algebra, if any, would be a manifestation of the Poisson brackets and not Dirac brackets. The noncommutative algebra was required to establish algebraic consistency of the boundary conditions with the basic Poisson brackets. For the free theory it was found that there was no clash between the boundary conditions and the Poisson brackets; hence there was no noncommutativity.

For the membrane interacting with a three-form potential a non-trivial algebraic relation was found that revealed the occurrence of noncommutativity, independent of any gauge choice or any approximations. Since this equation could not be solved, we passed on to its low-energy limit. Now this limit, which takes a membrane to a string, has been known for quite some time [18] and has been studied or exploited in several circumstances [17, 19, 20]. The cylindrical membrane is assumed to wrap around a circle, whose radius is taken to be vanishingly small. This enforces a double dimensional reduction with the

eleven-dimensional compactified target space passing over to the ten-dimensional space while the membrane effectively reduces to an open string. We have studied this limit in some detail and showed how the membrane boundary conditions, action and the world-volume metric were transformed into the corresponding expressions for the string. The equation governing noncommutativity in the membrane was likewise shown to reduce to the string example. Since every point in D-brane can correspond to the end points of the cylindrical membrane, we get noncommutativity in D-brane coordinates also — albeit in this low-energy limit. Of course, this feature of noncommutativity will persist even if this limit is not considered, otherwise the basic equation (90) becomes inconsistent.

It might be worthwhile to pursue the connection of the low-energy limit with the noncommutativity. Indeed, instead of taking a vanishingly small radius, it is possible to retain terms up to some orders of the radius [17]. This would presumably illuminate the nature of noncommutativity in the membrane (where it cannot be computed exactly) vis-à-vis the string.

Acknowledgements

The authors thank Pradip Mukherjee for discussions. BC and KK gratefully acknowledge help from a stimulating discussion with Koushik Ray. RB thanks the Japan Society for Promotion of Science (JSPS) for support and the members of the theory group, KEK, for their gracious hospitality. KK thanks the Council of Scientific and Industrial Research (CSIR), Government of India, for financial support.

Appendix: Noncommutativity in Open String

For an easy comparison of our results of open membrane with those of open string, we summarize here the essential results of [11].

The Free Polyakov string

The free Polyakov string action is

$$S_P = -\frac{T_S}{2} \int d\tau d\sigma \sqrt{-g} g^{ij} \partial_i X^\mu \partial_j X_\mu, \quad i, j = 0, 1, \quad (97)$$

where T_S stands for string tension and g_{ij} , up to a Weyl factor, is the induced metric $h_{ij} = \partial_i X^\mu \partial_j X_\mu$ on the world-sheet. The canonical momenta are

$$\Pi_\mu = -T_S \sqrt{-g} \partial^0 X_\mu, \quad \pi_{ij} = 0. \quad (98)$$

It is clear that while Π_μ are genuine momenta, $\pi_{ij} \approx 0$ are the primary constraints of the theory. To determine the secondary constraints, one can either follow the traditional Dirac's Hamiltonian approach, or just read it off from the equation obtained by varying g_{ij} since this is basically a Lagrange multiplier. This imposes the vanishing of the symmetric energy-momentum tensor:

$$T_{ij} = \frac{2}{\sqrt{-g}} \frac{\delta S_P}{\delta g^{ij}} = -T_S \partial_i X^\mu \partial_j X_\mu + \frac{T_S}{2} g_{ij} g^{kl} \partial_k X^\mu \partial_l X_\mu \approx 0. \quad (99)$$

Because of the Weyl invariance, the energy-momentum tensor is traceless:

$$T^i_i = g^{ij}T_{ij} = 0 \quad (100)$$

so that only two components of T_{ij} are independent. These components, which are the constraints of the theory, are given by

$$\chi_1 \equiv 2T_S g T^{00} = -2T_S T_{11} = \Pi^2 + T_S^2 h_{11} \approx 0, \quad (101)$$

$$\chi_2 \equiv \sqrt{-g} T^0_1 = \Pi_\mu \partial_1 X^\mu \approx 0. \quad (102)$$

The canonical Hamiltonian density obtained from (97) by a Legendre transformation is given by

$$\mathcal{H}_C = \sqrt{-g} T^0_0 = \frac{\sqrt{-g}}{2T_S g_{11}} \chi_1 + \frac{g_{01}}{g_{11}} \chi_2, \quad (103)$$

which, as expected, turns out to be a linear combination of the constraints. The boundary condition written in terms of phase-space variables is given by

$$\left[T_S \partial_1 X^\mu + \sqrt{-g} g^{01} \Pi^\mu \right]_{\sigma=0,\pi} = 0, \quad (104)$$

where the string parameters are in the region $-\infty \leq \tau \leq +\infty$, $0 \leq \sigma \leq \pi$. This boundary condition is incompatible with the first of the basic Poisson brackets:

$$\{X^\mu(\tau, \sigma), \Pi_\nu(\tau, \sigma')\} = \delta^\mu_\nu \delta(\sigma - \sigma'), \quad (105)$$

$$\{g_{ij}(\tau, \sigma), \pi^{kl}(\tau, \sigma')\} = \frac{1}{2} (\delta_i^k \delta_j^l + \delta_i^l \delta_j^k) \delta(\sigma - \sigma'). \quad (106)$$

From the basic Poisson brackets, it is easy to generate a first-class (involutive) algebra:

$$\begin{aligned} \{\chi_1(\sigma), \chi_1(\sigma')\} &= 4T_S^2 [\chi_2(\sigma) + \chi_2(\sigma')] \partial_1 \delta(\sigma - \sigma'), \\ \{\chi_2(\sigma), \chi_1(\sigma')\} &= [\chi_1(\sigma) + \chi_1(\sigma')] \partial_1 \delta(\sigma - \sigma'), \\ \{\chi_2(\sigma), \chi_2(\sigma')\} &= [\chi_2(\sigma) + \chi_2(\sigma')] \partial_1 \delta(\sigma - \sigma'). \end{aligned} \quad (107)$$

The constraints χ_1 and χ_2 generate the diffeomorphism transformations.

The boundary condition (104) is not a constraint in the Dirac sense, since it is applicable only at the boundary. Thus, there has to be an appropriate modification in the Poisson brackets, to incorporate this condition. This is not unexpected and occurs, for instance, in the example of a free scalar field $\phi(x)$ in $1+1$ dimensions, subjected to periodic boundary condition of period, say, 2π . There the Poisson bracket between the field $\phi(t, x)$ and its conjugate momentum $\pi(t, x)$ is given by

$$\{\phi(t, x), \pi(t, y)\} = \delta_P(x - y), \quad (108)$$

which is obtained automatically if one starts with the canonical harmonic-oscillator algebra for each mode in the Fourier space.

Before discussing the mixed condition (104), consider the simpler Neumann-type condition. Since the string coordinates $X^\mu(\tau, \sigma)$ transform as a world-sheet scalar under its reparametrization, it is more convenient to get back to scalar field $\phi(t, x)$ defined on $(1+1)$ -dimensional space-time, but with the periodic boundary condition replaced by Neumann boundary condition,

$$\partial_x \phi|_{\sigma=0,\pi} = 0, \quad (109)$$

at the end points of a 1-dimensional box of compact size, i.e. of length π . Correspondingly, the δ_P appearing there in the Poisson bracket (108) — consistent with periodic boundary condition — has to be replaced now with a suitable “delta function” incorporating Neumann boundary condition, rather than periodic boundary condition.

Note that the following usual property of delta function is also satisfied by $\delta_P(x - x')$:

$$\int_{-\pi}^{+\pi} dx' \delta_P(x - x') f(x') = f(x) \quad (110)$$

for any periodic function $f(x) = f(x + 2\pi)$ defined in the interval $[-\pi, +\pi]$. Restricting to the case of even and odd functions, $f_{\pm}(-x) = \pm f_{\pm}(x)$, the above integral reduces to

$$\int_0^{\pi} dx' \Delta_{\pm}(x, x') f_{\pm}(x') = f_{\pm}(x). \quad (111)$$

Since any function $\phi(x)$ defined in the interval $[0, \pi]$ can be regarded as a part of an even or odd function $f_{\pm}(x)$ defined in the interval $[-\pi, \pi]$, both $\Delta_{\pm}(x, x')$ act as delta functions defined in half of the interval at the right i.e. $[0, \pi]$ as follows from Eq. (111). It is still not clear which of these $\Delta_{\pm}(x, x')$ functions should replace $\delta_P(x - x')$ in the Poisson-bracket relation. Now consider the Fourier decomposition of an arbitrary function $f(x)$ satisfying periodic boundary condition $f(x) = f(x + 2\pi)$:

$$f(x) = \sum_{n \in \mathbb{Z}} f_n e^{inx}. \quad (112)$$

Clearly,

$$f'(0) = i \sum_{n>0} n(f_n - f_{-n}), \quad f'(\pi) = i \sum_{n>0} (-1)^n n(f_n - f_{-n}).$$

Now for even and odd functions, the Fourier coefficients are related as $f_{-n} = \pm f_n$ so that Neumann boundary condition $f'(0) = f'(\pi) = 0$ is satisfied if and only if $f(x)$ is even. Therefore, one has to regard the scalar field $\phi(x)$ defined in the interval $[0, \pi]$ and subjected to Neumann boundary condition (109) as a part of an even periodic function $f_+(x)$ defined in the extended interval $[-\pi, +\pi]$. It thus follows that the appropriate Poisson bracket for the scalar theory is given by

$$\{\phi(t, x), \pi(t, x')\} = \Delta_+(x, x').$$

It is straightforward to generalize it to the string case:

$$\{X^{\mu}(\tau, \sigma), \Pi_{\nu}(\tau, \sigma')\} = \delta_{\nu}^{\mu} \Delta_+(\sigma, \sigma'), \quad (113)$$

the Lorentz indices playing the role of “isospin” indices, as viewed from the world-sheet. Observe also that the other brackets

$$\{X^{\mu}(\tau, \sigma), X^{\nu}(\tau, \sigma')\} = 0, \quad (114)$$

$$\{\Pi^{\mu}(\tau, \sigma), \Pi^{\nu}(\tau, \sigma')\} = 0 \quad (115)$$

are consistent with the boundary conditions and hence remain unchanged.

The mixed condition (104) is compatible with the modified brackets (113) and (115), but not with (114). Therefore, make an ansatz:

$$\{X^{\mu}(\tau, \sigma), X^{\nu}(\tau, \sigma')\} = C^{\mu\nu}(\sigma, \sigma'), \quad (116)$$

where

$$C^{\mu\nu}(\sigma, \sigma') = -C^{\nu\mu}(\sigma', \sigma). \quad (117)$$

Imposing the boundary condition (104) on this algebra, one gets

$$\partial_1 C^{\mu\nu}(\sigma, \sigma')|_{\sigma'=0, \pi} = \partial_1 C^{\mu\nu}(\sigma, \sigma')|_{\sigma=0, \pi} = \sqrt{-g}g^{01}\eta^{\mu\nu}\Delta_+(\sigma, \sigma'). \quad (118)$$

For a restricted class of metrics that satisfy $\partial_1 g_{ij} = 0$ it is possible to give a quick solution of this equation as

$$C^{\mu\nu}(\sigma, \sigma') = \sqrt{-g}g^{01}\eta^{\mu\nu} [\Theta(\sigma, \sigma') - \Theta(\sigma', \sigma)]. \quad (119)$$

This noncommutativity can be made to vanish in gauges like conformal gauge, where $g^{01} = 0$, thereby restoring the usual commutative structure. The essential structure of the involutive algebra (107) is still preserved, only that $\delta(\sigma - \sigma')$ has to be replaced by $\Delta_+(\sigma, \sigma')$.

The Interacting Polyakov String

The Polyakov action for a bosonic string moving in the presence of a constant background Neveu-Schwarz two-form field $B_{\mu\nu}$ is given by

$$S_P = -\frac{T_S}{2} \int d\tau d\sigma \left[\sqrt{-g}g^{ij}\partial_i X^\mu \partial_j X_\mu + e\epsilon^{ij}B_{\mu\nu}\partial_i X^\mu \partial_j X^\nu \right], \quad (120)$$

where a ‘coupling constant’ e has been introduced. A usual canonical analysis leads to the following set of primary first-class constraints:

$$gT^{00} = \frac{1}{2} \left[(\Pi_\mu + eB_{\mu\nu}\partial_1 X^\nu)(\Pi^\mu + eB^{\mu\nu}\partial_1 X_\nu) + T_S^2 h_{11} \right] \approx 0, \quad (121)$$

$$\sqrt{-g}T^0{}_1 = \Pi_\mu \cdot \partial_1 X^\mu \approx 0, \quad (122)$$

where

$$\Pi_\mu = -T_S \left[\sqrt{-g}\partial^0 X_\mu + eB_{\mu\nu}\partial_1 X^\nu \right] \quad (123)$$

is the momentum conjugate to X^μ . The boundary condition written in terms of phase-space variables is

$$\left[\partial_1 X_\mu + \Pi^\rho (NM^{-1})_{\rho\mu} \right]_{\sigma=0, \pi} = 0, \quad (124)$$

where

$$M^\rho{}_\mu = T_S \left[\delta^\rho{}_\mu - e^2 B^{\rho\nu} B_{\nu\mu} \right], \quad N_{\nu\mu} = \frac{g_{01}}{\sqrt{-g}} \eta_{\nu\mu} + eB_{\nu\mu}. \quad (125)$$

The $\{X^\mu, \Pi_\nu\}$ Poisson bracket is the same as that of the free string whereas considering the general structure (116) and exploiting the above boundary condition, one obtains

$$\partial_1 C_{\mu\nu}(\sigma, \sigma')|_{\sigma=0, \pi} = (NM^{-1})_{\nu\mu} \Delta_+(\sigma, \sigma')|_{\sigma=0, \pi}. \quad (126)$$

As in the free case, restricting to the class of metrics satisfying $\partial_1 g_{ij} = 0$, the above equation has a solution

$$C_{\mu\nu}(\sigma, \sigma') = \frac{1}{2} (NM^{-1})_{(\nu\mu)} [\Theta(\sigma, \sigma') - \Theta(\sigma', \sigma)] + \frac{1}{2} (NM^{-1})_{[\nu\mu]} [\Theta(\sigma, \sigma') + \Theta(\sigma', \sigma) - 1]. \quad (127)$$

The modified algebra is gauge dependent; depends on the choice of the metric. However, there is no choice for which the noncommutativity vanishes. To show this, note that the origin of the noncommutativity is the presence of non-vanishing $N_{\nu\mu}$ in the boundary condition (124). Vanishing $N_{\nu\mu}$ would make $B_{\mu\nu}$ and $\eta_{\mu\nu}$ proportional which obviously cannot happen, as the former is an antisymmetric and the latter is a symmetric tensor. Hence, noncommutativity will persist for any choice of world-sheet metric g_{ij} . Specially interesting are the expressions for noncommutativity at the boundaries:

$$C_{\mu\nu}(0,0) = -C_{\mu\nu}(\pi,\pi) = -\frac{1}{2}(NM^{-1})_{[\nu\mu]}, C_{\mu\nu}(0,\pi) = -C_{\mu\nu}(\pi,0) = -\frac{1}{2}(NM^{-1})_{(\nu\mu)}. \quad (128)$$

References

- [1] For a review of the theory of membranes, see Jens Hoppe, [hep-th/0206192](#) and [13].
- [2] N. Seiberg and E. Witten, JHEP **9909** (1999) 032 [[hep-th/9908142](#)].
- [3] E. Bergshoeff, D. S. Berman, J. P. van der Schaar and P. Sundell, Nucl. Phys. **B590** (2000) 173 [[hep-th/0005026](#)].
- [4] F. Ardalan, H. Arfaei and M. M. Sheikh-Jabbari, JHEP **9902** (1999) 016 [[hep-th/9810072](#)].
- [5] F. Ardalan, H. Arfaei and M. M. Sheikh-Jabbari, Nucl. Phys. **B576** (2000) 578 [[hep-th/9906161](#)].
- [6] S. Kawamoto and N. Sasakura, JHEP **0007** (2000) 014 [[hep-th/0005123](#)].
- [7] A. Das, J. Maharana and A. Melikyan, JHEP **0104** (2001) 016 [[hep-th/0103229](#)].
- [8] Ken-Ichi Tezuka, Eur. Phys. J. **C25** (2002) 465 [[hep-th/0201171](#)].
- [9] Chong-Sun Chu and Pei-Ming Ho, Nucl. Phys. **B550** (1999) 151 [[hep-th/9812219](#)].
- [10] J. M. Romero and J. D. Vergara, [hep-th/0212035](#).
- [11] R. Banerjee, B. Chakraborty and S. Ghosh, Phys. Lett. **B537** (2002) 340 [[hep-th/0203199](#)].
- [12] A. Hanson, T. Regge and C. Teitelboim, *Constrained Hamiltonian Systems*, Roma, Accademia Nazionale Dei Lincei (1976).
- [13] W. Taylor, Rev. Mod. Phys. **73** (2001) 419 [[hep-th/0101126](#)].
- [14] S. Mandelstam, Phys. Rept. **13** (1974) 259.
- [15] C. Rebbi, Phys. Rept. **12** (1974) 1.
- [16] D. M. Gitman and I. V. Tyutin, *Quantization of Fields with Constraints*, Springer-Verlag, 1990.

- [17] U. Lindstrom, Phys. Lett. **B218** (1989) 315.
- [18] M. J. Duff, Paul S. Howe, T. Inami and K. S. Stelle, Phys. Lett. **B191** (1987) 70.
- [19] E. Bergshoeff, E. Sezgin and P. K. Townsend, Annals Phys. **185** (1988) 330.
- [20] L. Smolin, Phys. Rev. **D57** (1998) 6216 [[hep-th/9710191](#)].